

The correspondence section is a public forum and, as such, is not peer-reviewed. EHP is not responsible for the accuracy, currency, or reliability of personal opinion expressed herein; it is the sole responsibility of the authors. EHP neither endorses nor disputes their published commentary.

Atmospheric Interactions and Cardiac Arrhythmias

<http://dx.doi.org/10.1289/ehp.1409636>

Although plausible pathophysiological mechanisms link air pollution to arrhythmogenesis, among them altered autonomic tone, repolarization abnormalities, oxidative stress, myocardial ischemia, and increased intracardiac pressure (Link and Dockery 2010), definitive conclusions have not been reached as yet. Langrish et al. (2014) analyzed 13 double-blind randomized crossover studies and found no significant risk of arrhythmia attributable to acute controlled exposure to air pollutants. Three issues related to meteorological factors probably either confound or modify the short-term association between air pollution and cardiac arrhythmia.

First, several meteorological elements, including air temperature, atmospheric pressure, relative air moisture, and wind speed and direction, also are implicated in triggering ventricular (Čulić et al. 2004, 2005) and supraventricular (Čulić et al. 2012, 2013) arrhythmias independent of physical and emotional stress. In the short term, those meteorological factors may facilitate arrhythmias in susceptible patients by increasing circulatory load and thromboinflammatory processes (Čulić 2014).

Second, these same meteorological elements substantially influence concentrations of sulfur dioxide, carbon monoxide, nitrogen dioxide, ozone, and suspended particulate matter (Bertaccini et al. 2012; Ilten and Selici 2008; Ito et al. 2007). In addition, the greatest ozone production and pollution results from stable, dry, hot weather with high atmospheric pressure and low wind (Vanos et al. 2014).

Air pollution may increase human vulnerability to the effects of temperature, and temperature extremes, in turn, influence population vulnerability to air pollution (Burkart et al. 2013; Ren et al. 2006). Vanos et al. (2014) reported that cardiovascular and respiratory mortality due to short-term exposure to gaseous air pollutants was significantly modified by weather types and season. Alberdi et al. (1998) reported that both relative air moisture and air temperature are strongly related to daily mortality even after controlling for air pollution and influenza. Keatinge and Donaldson (2001) suggested that prolonged cold weather with less wind and rain may produce false associations between mortality and certain air pollutants.

Finally, strong mutual interrelations exist among the above-mentioned meteorological

elements. Alberdi et al. (1998) pointed out the strong inverse association they observed between relative air moisture and air temperature as an important problem for regression analysis.

Langrish et al. (2014) caution against definitive acceptance of air pollution as an independent trigger of cardiac arrhythmias. However, the studies included in their analysis had no data on meteorological factors. It is likely that interactive effects among air pollutants and meteorological elements bias each other's association with arrhythmias and other acute cardiac events. Therefore, further research of the health effects of atmospheric factors should continue in order to identify potentially harmful influences for the population as whole as well as for its vulnerable subgroups.

The author declares he has no actual or potential competing financial interests.

Viktor Čulić

Department of Cardiology, University Hospital Center Split, Split, Croatia
E-mail: viktor.culic@st.t-com.hr

REFERENCES

- Alberdi JC, Díaz J, Montero JC, Mirón I. 1998. Daily mortality in Madrid community 1986–1992: relationship with meteorological variables. *Eur J Epidemiol* 14(6):571–578; PMID:9794124.
- Bertaccini P, Dukić V, Ignaccolo R. 2012. Modeling the short-term effect of traffic and meteorology on air pollution in Turin with generalized additive models. *Adv Meteorol*, article ID 609328; doi:10.1155/2012/609328.
- Burkart K, Canário P, Breiten S, Schneider A, Scherber K, Andrade H, et al. 2013. Interactive short-term effects of equivalent temperature and air pollution on human mortality in Berlin and Lisbon. *Environ Pollut* 183:54–63; doi:10.1016/j.envpol.2013.06.002.
- Čulić V, Eterović D, Mirić D, Giunio L, Lukin A, Fabijanić D. 2004. Triggering of ventricular tachycardia by meteorologic and emotional stress: protective effect of β -blockers and anxiolytics in men and elderly. *Am J Epidemiol* 160(11):1047–1058; doi:10.1093/aje/kwh335.
- Čulić V, Silić N, Mirić D. 2005. Triggering of ventricular ectopic beats by emotional, physical and meteorologic stress: role of age, sex, medications, and chronic risk factors. *Croat Med J* 46(6):894–906; PMID:16342342.
- Čulić V, Silić N, Mirić D. 2012. Triggering of supraventricular premature beats. The impact of acute and chronic risk factors. *Int J Cardiol* 158(1):112–117; doi:10.1016/j.ijcard.2012.04.059.
- Čulić V, Silić N, Hodžić M. 2013. Triggering of supraventricular tachycardia by physical activity and meteorologic factors. *Int J Cardiol* 168(4):4295–4300; doi:10.1016/j.ijcard.2013.04.195.
- Čulić V. 2014. Inflammation, coagulation, weather and arrhythmogenesis: is there a linkage? *Int J Cardiol* 176(1):289–293; doi:10.1016/j.ijcard.2014.06.078.
- Ilten N, Selici AT. 2008. Investigating the impacts of some meteorological parameters on air pollution in Balikesir, Turkey. *Environ Monit Assess* 140(1–3):267–277; doi:10.1007/s10661-007-9865-1.
- Ito K, Thurston GD, Silverman RA. 2007. Characterization of PM_{2.5}, gaseous pollutants, and meteorological interactions in the context of time-series health effects models. *J Exp Sci Environ Epidemiol* 17(suppl 2):S45–S60; doi:10.1038/sj.jes.7500627.

- Keatinge W, Donaldson G. 2001. Mortality related to cold and air pollution in London after allowance for effects of associated weather patterns. *Environ Res* 86(3):209–216; doi:10.1006/enrs.2001.4255.
- Langrish JP, Watts SJ, Hunter AJ, Shah ASV, Bosson JA, Unosson J, et al. 2014. Controlled exposures to air pollutants and risk of cardiac arrhythmia. *Environ Health Perspect* 122(7):747–753; doi:10.1289/ehp.1307337.
- Link MS, Dockery DW. 2010. Air pollution and the triggering of cardiac arrhythmias. *Curr Opin Cardiol* 25(1):16–22; doi:10.1097/HCO.0b013e32833358cd.
- Ren C, Williams GM, Tong S. 2006. Does particulate matter modify the association between temperature and cardiorespiratory diseases? *Environ Health Perspect* 114(11):1690–1696; doi:10.1289/ehp.9266.
- Vanos JK, Hebbert C, Cakmak S. 2014. Risk assessment for cardiovascular and respiratory mortality due to air pollution and synoptic meteorology in 10 Canadian cities. *Environ Pollut* 185:322–332; doi:10.1016/j.envpol.2013.11.007.

Atmospheric Interactions and Cardiac Arrhythmias: Langrish et al. Respond

<http://dx.doi.org/10.1289/ehp.1409636R>

We agree with Čulić's argument (2015) that environmental influences on human health are complex and likely multifactorial. Exposure to indoor and ambient urban air pollution has been estimated to contribute to 7 million premature deaths each year, predominantly from cardiovascular and respiratory conditions (Lim et al. 2012). Associations between exposure and cardiovascular mortality and morbidity have been demonstrated for nitrogen dioxide, sulfur dioxide, ozone, carbon monoxide, and particulate matter (Brook et al. 2010), although the associations are strongest for fine and ultrafine particulate matter (Hoek et al. 2013).

Air pollution is extremely complex and consists not of single components in isolation but rather combinations of components. These constituent components interact with one another in the environment, which may alter potential toxicity and the subsequent health impacts. Meteorological factors such as wind speed and direction, humidity, atmospheric pressure, and temperature play an important part both in determining an individual's exposure to ambient air pollution and in affecting the concentrations, chemical composition, and clearance of elements of the air pollution mixture. This is particularly true for the secondary pollutant ozone, which has a strong relationship with season and temperature (Langrish et al. 2010b). We agree that, in assessing the impact of ambient air pollution on public health, it is important to assess the air pollution mixture as a whole.

In our studies we used a controlled exposure facility to assess, in a robust and well-validated fashion (Langrish et al. 2010a), the contribution to potential arrhythmogenesis of individual air pollutants—diesel exhaust, wood smoke, ozone, and nitrogen dioxide—as well as ambient air pollution in Beijing,

China (Langrish et al. 2014). Exposure to neither air pollutants in isolation nor ambient Beijing air pollution was associated with cardiac dysrhythmia in either patients with coronary heart disease or healthy volunteers. As such, our studies do not address the influence of meteorological conditions on an individual's risk of cardiac arrhythmia; indeed, the meteorological conditions in Beijing were fairly constant throughout our studies (Langrish et al. 2009, 2012).

There is emerging evidence that cardiovascular morbidity and mortality is associated with meteorological and environmental conditions, and we agree with Čulić's statement that further research on the health impacts of atmospheric factors is important both for public health and for better understanding the interaction between urban air pollution and external influences.

The authors declare they have no actual or potential competing financial interests.

Jeremy P. Langrish,¹ Jenny A. Bosson,² Thomas Sandström,² Anders Blomberg,² David E. Newby,¹ and Nicholas L. Mills¹

¹University/BHF Centre for Cardiovascular Science, University of Edinburgh, United Kingdom; ²Department of Public Health and Clinical Medicine, Division of Medicine/Respiratory Medicine, Umeå University, Umeå, Sweden
E-mail: d.e.newby@ed.ac.uk

REFERENCES

- Brook RD, Rajagopalan S, Pope CA III, Brook JR, Bhatnagar A, Diez-Roux AV, et al. 2010. Particulate matter air pollution and cardiovascular disease. An update to the scientific statement from the American Heart Association. *Circulation* 121:2331–2378; doi:10.1161/CIR.0b013e3181d8ce1.
- Čulić V. 2015. Atmospheric interactions and cardiac arrhythmias. *Environ Health Perspect* 123(6):A144; doi:10.1289/ehp.1409636.
- Hoek G, Krishnan RM, Beelen R, Peters A, Ostro B, Brunekreef B, et al. 2013. Long-term air pollution exposure and cardio-respiratory mortality: a review. *Environ Health* 12:43; doi:10.1186/1476-069X-12-43.
- Langrish JP, Mills NL, Chan JK, Leseman DL, Aitken RJ, Fokkens PH, et al. 2009. Beneficial cardiovascular effects of reducing exposure to particulate air pollution with a simple facemask. *Part Fibre Toxicol* 6:8; doi:10.1186/1743-8977-6-8.
- Langrish JP, Frampton M, Blomberg A. 2010a. Human exposure studies. In: *Cardiovascular Effects of Inhaled Ultrafine and Nanosized Particles* (Casseo F, Mills N, Newby D, eds). Hoboken, NJ: John Wiley & Sons, 217–239.
- Langrish JP, Mills NL, Donaldson K, Newby DE. 2010b. Response to Peter Joseph. *Heart* 96(6):472–473; doi:10.1136/hrt.2009.189308.
- Langrish JP, Li X, Wang S, Lee MM, Barnes GD, Miller MR, et al. 2012. Reducing personal exposure to particulate air pollution improves cardiovascular health in patients with coronary heart disease. *Environ Health Perspect* 120(3):367–372; doi:10.1289/ehp.1103898.
- Langrish JP, Watts SJ, Hunter AJ, Shah AS, Bosson JA, Unosson J, et al. 2014. Controlled exposures to air pollutants and risk of cardiac arrhythmia. *Environ Health Perspect* 122(7):747–753; doi:10.1289/ehp.1307337.
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. 2012. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380(9859):2224–2260; doi:10.1016/S0140-6736(12)61766-8.